
**DEPARTMENT OF THE INTERIOR
U.S. GEOLOGICAL SURVEY**

**Pliocene planktic foraminifer census data from the
North Atlantic region**

PRISM Project Members
US Geological Survey, Reston, VA 20192



Open-File Report 96-669

This report is preliminary and has not been reviewed for conformity with
U.S. Geological Survey editorial standards.

Pliocene planktic foraminifer census data from the North Atlantic region

PRISM Project Members

U.S. Geological Survey, 955 National Center, Reston, VA 20192

INTRODUCTION

The U.S. Geological Survey is conducting a long-term study of the climatic and oceanographic conditions of the Pliocene known as PRISM (Pliocene Research, Interpretation, and Synoptic Mapping). One of the major elements of the study involves the use of quantitative composition of planktic foraminifer assemblages to estimate sea-surface temperatures and identify major oceanographic boundaries and water masses (Dowsett, 1991; Dowsett and Poore, 1991; Dowsett et al., 1992; Dowsett et al., 1994). We have analyzed more than 900 samples from 19 core sites in the North Atlantic Basin (Fig. 1) resulting in a large volume of raw census data. These data are presented here together to facilitate comparison of North Atlantic faunal assemblages.




Figure 1. Location of sites discussed in text.

Latitude, longitude, water depth, source of faunal data and source of data used to construct age model (or publication from which age model was taken) are provided for each locality in Table 1. All ages refer to the geomagnetic polarity time scale of Berggren et al. (1985). Counts of species tabulated in each sample are given in Tables 2-20. DSDP and ODP sample designations are abbreviated in Tables 2-20 as core-section, depth within section in centimeters (eg. 10-5, 34 = core 10, section 5, 34 cm below top of section 5).

Table 1. Location and water depth of sites included in this report, original sources of faunal data, and reference for age models included with faunal data sets.

Site	Latitude	Longitude	Water depth (m)	Original faunal data source	Age model
366A	5.68	-19.85	2860	Dowsett et al., 1988	Cepek et al., 1978
396	22.90	-43.50	4450	Wiggs & Dowsett, 1992	Bukry, 1978
410	45.51	-29.48	2975	Dowsett et al., 1988	Poore, 1978
502A	11.49	-79.38	3051	Wiggs & Poore, 1991	Dowsett, 1989
541	15.50	-58.73	4940	Dowsett & Polanco, 1992	Wilson, 1984
546	33.80	-9.60	3958	Dowsett & Polanco, 1992	Hinz et al., 1984
548	48.50	-12.00	1251	Loubere & Moss, 1986	Dowsett & Loubere, 1992
552A	56.04	-23.23	2301	Dowsett & Poore, 1990	Dowsett & Poore, 1990
603C	35.49	-70.03	4633	Poore, 1991	Canninga et al., 1986
606	37.34	-35.50	3007	Dowsett et al., 1988	Dowsett, 1989
607	41.00	-32.96	3427	Dowsett & West, 1992	Baldauf et al., 1986
608	42.90	-23.10	3526	This paper	Baldauf et al., 1986
609B	49.88	-24.24	3883	This paper	Baldauf et al., 1986
610A	53.22	-18.89	2417	This paper	Baldauf et al., 1986
646B	58.25	-48.33	3451	Dowsett et al., 1988	Baldauf et al., 1989
659A	18.00	-21.10	3070	Foley & Dowsett, 1992	Weaver et al., 1989
661A*	9.45	-19.39	4006	Dowsett & West, 1992	Weaver et al., 1989
667A	4.55	-21.90	3529	Foley and Dowsett, 1992	Weaver et al., 1989
672A	15.50	-58.50	4975	Wiggs and Dowsett, 1992	Moore, Mascle, et al., 1988

*R.Z. Poore responsible for faunal data

METHODS

The samples used in this study were processed using low temperature (isotopic) techniques. This includes sample disaggregation in water, drying at 50°C, and sieving into 63 µm - 149 µm and 149 µm size fractions. Following standard procedures, some samples required additional treatment with Calgon or H₂O₂ added to the wash to obtain clean specimens.

A split of 300-350 planktic foraminifer specimens was obtained from the 149 µm size fraction using a Carpc sample splitter. Specimens were identified, sorted, and fixed to a standard 60-square micropaleontological slide.

COUNTING CATEGORIES

Taxa included in Tables 2-20 are summarized in a comprehensive list below. In general, our taxonomic concepts follow Parker (1962; 1967) and Blow (1969); exceptions to their practices are noted below.

Candeina nitida d'Orbigny

Dentoglobigerina altispira (Cushman and Jarvis)

Globoquadrina venezuelana (Hedberg)

Globigerina bulloides (d'Orbigny)
We include *Globigerina praebulloides* Blow in this category.

Globigerina calida Parker
Globigerina conglomerata (Schwager)

Globigerina decoraperta Takayanagi and Saito

Globigerina digitata Brady

Globigerina eamesi Blow

Globigerina falconensis Blow

Globigerina incisa (Bronnimann and Resig)

Globigerina nepenthes Todd

Globigerina praedigitata Parker

Globigerina pseudobesa (Salvatorini)

Globigerina pseudobulloides (Plummer)

Globigerina sp. 1
Taxon resembles *G. falconensis* but has reticulate surface texture similar to *G. woodi* group.

Globigerina woodi Jenkins
We include specimens of *Globigerina apertura* Cushman in this category.

Globigerinella aequilateralis (Brady)

Globigerinella siphonifera (d'Orbigny)

Globigerinita glutinata (Egger)

Globigerinoides conglobatus (Brady)

Globigerinoides obliquus Bolli
We include *G. extremus* Bolli and Bermudez in this category.

Globigerinoides ruber (d'Orbigny)

Globigerinoides sacculifer (Brady) s.l.
This category includes *G. quadrilobatus* (d'Orbigny) and *G. trilobus* (Reuss).

Globigerinoides spp.
Representatives of *Globigerinoides* (usually small) that could not be confidently assigned to *G. ruber*, *G.*

obliquus (s.l.) or *G. conglobatus* are included in this category.

Globorotalia cibaoensis Bermudez

Globorotalia conomiozea Kennett

Globorotalia crassaformis (Galloway and Wissler) For the purpose of this report we have combined the census data of a number of taxa including *G. ronda* Blow, *G. oceanica* Cushman and Bermudez, *G. viola* Blow, and *G. crassula* Blow into this category.

Globorotalia hirsuta (d'Orbigny)

Globorotalia margaritae Bolli and Bermudez

Globorotalia menardii (Parker, Jones, and Brady) This category includes various members of the *G. menardii* lineage such as *G. limbata* (Fornasini) and *G. miocenica* Palmer.

Globorotalia praepumilio (Parker)

Globorotalia pumilio Parker
This category includes small forms with 5-7 chambers in the ultimate whorl that are similar to *Globorotalia pumilio* Parker, *G. praepumilio* (Parker) and *G. pseudopumilio* Bronnimann and Resig.

Globorotalia puncticulata (Deshayes) s.l.
This category includes *G. inflata* (d'Orbigny).

Globorotalia scitula (Brady) s.l.
This category includes various members of the *G. scitula* group, for example *G. subscitula* Conato.

Globorotalia spp.
This category includes rare *Globorotalia* not assigned to the other taxa listed here.

Globorotalia tosaensis Takayanagi and Saito We include occurrences of

Globorotalia truncatulinoides
(d'Orbigny) in this category.

Globorotalia tumida (Brady) s.l.
This category includes *G. plesiotumida*
Blow and Banner.

Globorotaloides hexagona (Natland)

Neogloboquadrina acostaensis (Blow)
We include *N. continuosa* (Blow) in this
category.

Neogloboquadrina atlantica (Berggren)
We separate sinistral and dextral coiled
specimens. See Poore and Berggren
(1975) for discussion of this highly
variable taxon.

Neogloboquadrina humerosa
(Takayanagi and Saito)

Neogloboquadrina pachyderma
(Ehrenberg) Sinistral variety includes
relatively small, compact
Neogloboquadrina with 4-5 chambers in
the ultimate whorl, kummerform ultimate
chamber, and a slightly to distinct oval
equatorial outline. Separating small
sinistral *N. atlantica* from large sinistral
N. pachyderma is arbitrary in many
North Atlantic high-latitude sites. Dextral
variety is confined to 4 chambers in the
final whorl. Dextral coiled specimens
with 4 1/2 chambers in the final whorl are
tabulated in a "dupac" category.

Neogloboquadrina spp.
This category includes
Neogloboquadrina that were not
identified to specific level but generally
does not include representatives of *N.*
atlantica.
Orbulina universa d'Orbigny

Pulleniatina obliquiloculata (Parker and
Jones)

Sphaeroidinellopsis spp.
This category includes members of the
genera *Sphaeroidinella* and
Sphaeroidinellopsis.

Turborotalita quinqueloba (Natland)

OTHER

This category includes unidentified
specimens and taxa that are rare within
assemblages from the cores.

TOTAL PLANKTICS

Total number of planktic forams in the
counting split

FRAGMENTS

Fragments of planktic foraminifers

BENTHICS

Number of benthic foraminifers in
planktic counting split

ACKNOWLEDGMENTS

This report and the census data it contains
are products of the PRISM (Pliocene
Research, Interpretation, and Synoptic
Mapping) Project. A large number of
scientists and technical and administrative
support staffers are due thanks for the
help they provided while these data were
being produced. We thank Tom Cronin,
Scott Ishman, and Debra Willard for
reviewing this manuscript and
acknowledge the technical help of Jean
Self-Trail and Marci Robinson in the
laboratory. We also thank the Ocean
Drilling Program for supplying samples
and thus making this study possible.

REFERENCES

- Baldauf, J.G., Aksu, A.E., et al., 1989. Magnetostratigraphic and biostratigraphic synthesis of Ocean Drilling Program Leg 105: Labrador Sea and Baffin Bay. *Proceedings of the Ocean Drilling Program, Scientific Results* 105: 935-956.
- Baldauf, J.G., Thomas, E., Clement, B., Takayama, T., Weaver, P.P.E., Backman, J., Jenkins, G., Mudie, P.J. and Westberg-Smith, M.J., 1986. Magnetostratigraphic and biostratigraphic synthesis, Deep Sea Drilling Project Leg 94. *Initial Reports of the Deep Sea Drilling Project* 94: 1159-1205.
- Berggren, W.A., Kent, D.V. and Van Couvering, J.A., 1985. Neogene geochronology and chronostratigraphy. *The Chronology of the Geological Record*. London, Blackwell Scientific Publications. 211-260.
- Blow, W. H., 1969, Late middle Eocene to Recent planktonic foraminiferal biostratigraphy. In Bronnimann, P. and Renz, H. H., (Eds), *Proceedings of First Planktonic Conference*: Leiden (E. J. Brill), p. 199-422.
- Bukry, D., 1977. Cenozoic coccolith and silicoflagellate stratigraphy, offshore northwest Africa, Leg 41, Deep Sea Drilling Project. *Initial Reports of the Deep Sea Drilling Project* 41: 689-708.
- Bukry, D., 1978. Neogene coccolith stratigraphy, mid-Atlantic Ridge, Deep Sea Drilling Project, Leg 45. *Initial Reports of the Deep Sea Drilling Project* 45: 307-312.
- Canninga, G., Zijderveld, J.D.A. and van Hinte, J.E., 1986. Late Cenozoic magnetostratigraphy of Deep Sea Drilling Project Hole 603C, Leg 93, On the North American Continental Rise of Cape Hatteras. *Initial Reports of the Deep Sea Drilling Project* 93: 839-848.
- Cepek, P., Johnson, D., Krasheninnikov, V., and Pflaumann, U., 1978. Synthesis of the Leg 41 biostratigraphy and paleontology, Deep Sea Drilling Project. *Initial Reports of the Deep Sea Drilling Project* 41: 1181-1198.
- Dowsett, H.J., 1989. Application of the graphic correlation method to Pliocene marine sequences. *Marine Micropaleontology* 14: 3-32.
- Dowsett, H. J., 1991, The development of a long-range foraminifer transfer function and application to Late Pleistocene North Atlantic climatic extremes. *Paleoceanography* 6: 259-273.
- Dowsett, H.J., Cronin, T.M., Poore, R.Z., Thompson, R.S., Whatley, R.C. and Wood, A.M., 1992. Micropaleontological evidence for increased meridional heat transport in the North Atlantic Ocean during the Pliocene. *Science* 258: 1133-1135.
- Dowsett, H.J., Gosnell, L.B. and Poore, R.Z., 1988. Pliocene planktic foraminifer census data from Deep Sea Drilling Project Holes 366A, 410, 606, and 646B. *U.S. Geological Survey Open File Report* 88-654: 14.
- Dowsett, H.J. and Loubere, P., 1992. High resolution Late Pliocene sea-surface temperature record from the

- Northeast Atlantic Ocean. *Marine Micropaleontology* 20: 91-105.
- Dowsett, H.J. and Polanco, E.F. 1992, Pliocene planktic foraminifer census data from Deep Sea Drilling Project Holes 541 and 546: *U.S. Geological Survey Open File Report* 92-418, 4p.
- Dowsett, H.J. and Poore, R.Z., 1990. A new planktic foraminifer transfer function for estimating Pliocene through Holocene Sea Surface temperatures. *Marine Micropaleontology* 16(1/2): 1-23.
- Dowsett, H. J. and R. Z. Poore, 1991, Pliocene sea surface temperatures of the North Atlantic Ocean at 3.0 Ma, *Quaternary Science Reviews* 10: 189-204.
- Dowsett, H.J. and West, S.M. 1992, Pliocene planktic foraminifer census data from Deep Sea Drilling Project Hole 607 and Ocean Drilling Program Hole 661A: *U.S. Geological Survey Open File Report* 92-413, 4p.
- Dowsett, H.J., Thompson, R.S., Barron, J.A., Cronin, T.M., Fleming, R.F., Ishman, S.E., Poore, R.Z., Willard, D.A. and Holtz, T.R., 1994. Joint investigations of the middle Pliocene climate I: PRISM paleoenviromental reconstructions. *Global and Planetary Change* 9(4): 169-195.
- Foley, K.M. and Dowsett, H.J., 1992. Pliocene planktic foraminifer census data from Ocean Drilling Program Holes 667 and 659A. *U.S. Geological Survey Open File Report* 92-434, 8p.
- Hinz, K., Winterer, E.L., et al., 1984. Site 546. *Initial Reports of the Deep Sea Drilling Project* 79: 179-221.
- Loubere, P. and Moss, K., 1986. Late Pliocene climatic change and the onset of Northern Hemisphere glaciation as recorded in the northeast Atlantic Ocean. *Geological Society of America Bulletin* 97: 818-828.
- Moore, J.C., Mascle, A., et al., 1988. Site 672. *Proceedings of the Ocean Drilling Program, Initial Reports* 110: 205-310.
- Parker, F. L., 1962, Planktonic foraminiferal species in Pacific sediments, *Micropaleontology* 8: 219-254.
- Parker, F. L., 1967, Late Tertiary biostratigraphy (Planktonic Foraminifera) of tropical Indo-Pacific deep-sea cores: *Bulletins of American Paleontology* 52: 115-208.
- Poore, R.Z., 1978. Oligocene through Quaternary planktonic foraminiferal biostratigraphy of the North Atlantic: DSDP Leg 49. *Initial Reports of the Deep Sea Drilling Project* 49: 447-517.
- Poore, R.Z., 1991. Pliocene planktic foraminifer census data from Deep Sea Drilling Project Hole 603C. *U.S. Geological Survey Open File Report* 91-309, 7p.
- Poore, R. Z., and Berggren, W. A., 1975, The morphology and classification of *Neoglobobulimina atlantica* (Berggren), *Journal of Foraminiferal Research* 5: 77-84.
- Weaver, P.P.E., Backman, J., et al., 1989. Biostratigraphic synthesis, Leg 108, Eastern Equatorial Atlantic. *Proceedings of the Ocean Drilling Program, Scientific Results* 108: 455-462.
- Wiggs, L.B. and Dowsett, H.J., 1992. Pliocene planktic foraminifer census

data from Deep Sea Drilling Project
Hole 396 and Ocean Drilling Program
Hole 672. *U.S. Geological Survey
Open File Report* : 92-414, 5p.
Wiggs, L.B. and Poore, R.Z., 1991.
Pliocene planktic foraminifer census
data from Deep Sea Drilling Project

Holes 502A, B, C. *U.S. Geological
Survey Open File Report* 91-325.

Wilson, D.S., 1984. Paleomagnetic
results from Deep Sea Drilling Project
Leg 78A. *Initial Reports of the Deep
Sea Drilling Project* 78: 583-591.

This report is available from USGS Information Services, Box 25286, Denver, Colorado,
80225 or <URL:<http://geochange.er.usgs.gov/pub/PRISM/>>.

For more information, contact:

Harry Dowsett
U.S.G.S.
955 National Center
Reston, VA 20192
(703) 648-5282
hdowsett@geochange.er.usgs.gov

Dick Poore
U.S.G.S.
955 National Center
Reston, VA 20192
(703) 648-5270
rpoore@geochange.er.usgs.gov